

Does Repeated Use of a Cereal Rye Cover Crop Reduce the Need for N Fertilizer for Corn?

In a Nutshell:

- Jack Boyer has been planting a cereal rye cover crop in a corn-soybean rotation on his farm for six years. Because Boyer knows that a cover crop can improve soils, he was curious if the N fertilizer rate he applies to corn could be reduced due to several years of repeated cover crop use. He compared his typical N rate (180 lb N/ac) with a reduced rate (130 lb N/ac).
- Boyer hypothesized that corn yields would be comparable between the two N fertilizer rates; as such, reducing the N rate would be more profitable due to lowered input costs.

Key Findings:

- Cornstalk nitrate concentration – which can indicate if the corn had a sufficient supply of N during the season – was not affected by N fertilizer rate.
- Boyer recorded equal corn yields between the reduced and typical N rates.
- Reducing the N rate by 50 lb N/ac improved Boyer's returns by \$26/ac.

METHODS

Design

Boyer planted corn on Apr. 23, 2019 at a population of 35,000 seeds/ac to a field that was planted to soybeans in 2018. Following soybean harvest in the previous fall, Boyer seeded a cereal rye cover crop to the field at a rate of 50 lb/ac on Oct. 23, 2018. He terminated the cover crop and controlled early-season weeds on May 4 with atrazine (0.25 lb/ac), Outlook (14 oz/ac), Durango (48 oz/ac) and Liberty (1 qt/ac). Boyer performed another weed control pass with Zaar (16 oz/ac) and Impact (9.5 oz/ac) on June 13.

To test whether he could reduce his N fertilizer rate, Boyer compared two treatments:

1. 130 lb N/ac (reduced rate);
2. 180 lb N/ac (typical rate).

Boyer implemented four replications of the two treatments in strips that measured 30 ft wide and 1,570 ft long (**Figure A1**). He applied 100 lb N/ac as anhydrous ammonia on Apr. 16 and 30 lb N/ac as UAN(28%) on May 4 to all strips. Boyer then applied 50 lb N/ac as sidedress on June 14 but only to the strips assigned to his typical N rate.

Measurements

On June 12, we collected soil samples to a 12-



Jack Boyer: "This trial was worthwhile because it showed I was able to reduce N input without reducing yield."

in. depth from several areas in the field. We sent these samples to Ag Source Laboratories (Ellsworth, IA) to determine the availability of soil nitrate to corn prior to Boyer sidedressing the strips receiving the typical N rate. This is commonly known as the late-spring soil nitrate test (LSNT).^[1]

Theo Gunther, from the Iowa Soybean Association, collected cornstalk samples from each strip on Oct. 4 and sent the samples to Midwest Labs (Omaha, NE). The lab determined the concentration of nitrate in the

stalks – this can be an indicator of sufficient, insufficient or excessive availability of N to the corn during the growing season.^[2]

On Oct. 26, Boyer harvested the corn and recorded yields from each individual strip.

Data analysis

To evaluate any effects of N rate, we calculated the least significant difference (LSD) at the 95% confidence level using a t-test for each measurement: cornstalk nitrate concentration and corn yield. For each measurement, if the

Cooperators

Jack Boyer – Reinbeck

Funding

USDA-NRCS
Walton Family Foundation

difference resulting from the two N rate treatments was *greater* than the LSD, we would expect such a difference to occur 95 times out of 100 – we refer to this as a statistically significant effect. On the other hand, if the difference resulting from the two N rate treatments was *less* than the LSD, we consider the results to be statistically similar. We could make these statistical calculations because Boyer’s experimental design involved replication of the two treatments (**Figure A1**).

RESULTS AND DISCUSSION

Late-spring soil nitrate test

Soil samples collected on June 12 revealed that Boyer’s field had an average nitrate-N concentration of 17 ppm. This test result would suggest the application of 64 lb N/ac as sidedress to Boyer’s corn. We used this test result purely as a reference – Boyer did not sidedress any N to the reduced rate treatment and he sidedressed 50 lb N/ac to the typical rate treatment per the design of this trial.

Cornstalk nitrate concentration

N fertilizer rate had no statistical effect on stalk nitrate concentrations (**Table 1**). Across the two N rates, the average stalk nitrate concentration was 408 ppm. This value suggests that Boyer’s corn (regardless of N rate) had access to the minimal amount of N necessary for maximum yield potential, according to ISU Extension and Outreach.^[2]

Corn yield and economic returns

Corn yield was not affected by N fertilizer rate (**Figure 1**). Recall that the results of the LSNT mentioned before recommended a sidedress application of 64 lb N/ac to Boyer’s corn in June. As such, we might have expected the sidedress application of 50 lb N/ac as part of Boyer’s typical rate (180 lb N/ac) to increase yield compared to the reduced rate (130 lb N/ac). But this was not the case in this trial as corn yields were statistically similar between the two N fertilizer rates.

Because Boyer saw similar yields between the two N rates, it is important to consider the differences in costs and returns between the two N rates. Boyer’s typical N rate (180 lb N/ac) involved a sidedress application of 50 lb N/ac as anhydrous ammonia on June 14. The 50 lb N/ac cost \$13.50/ac, according to Boyer, and the sidedress application cost \$12.50/ac, according to ISU’s survey of farm custom rates in 2019.^[3] This amounted to \$26/ac in additional costs compared to the reduced rate which did not receive a sidedress application. Therefore, because Boyer saw similar yields between the two N rates (**Figure 1**), his economic returns were improved by \$26/ac with the reduced rate (130 lb N/ac) compared with his typical rate (180 lb N/ac).

CONCLUSIONS AND NEXT STEPS

Boyer conducted this study because he was curious about opportunities to reduce input costs and either maintain or improve profitability. Just as

TABLE 1. Stalk nitrate concentrations at Jack Boyer’s in 2019.

TREATMENT	STALK NITRATE CONCENTRATION (ppm)
130 lb N/ac (reduced rate)	600
180 lb N/ac (typical rate)	215
<i>Trt difference</i>	385
<i>LSD (95%)</i>	805
Because the difference between the treatment means is less than the least significant difference (LSD), the treatments are considered statistically similar at the 95% confidence level.	

Corn yields




FIGURE 1. Corn yields at Jack Boyer’s, harvested on Oct. 26, 2019. Columns represent yields for each individual strip. Above each set of columns is the mean for both N fertilizer rate treatments. Because the difference between the two means (6 bu/ac) is less than the least significant difference (LSD; 10 bu/ac), the treatments are considered statistically similar at the 95% confidence level.

Boyer had suspected, he was able to maintain corn yields while reducing his N fertilizer rate by 50 lb N/ac. In this case, applying 130 compared to 180 lb N/ac reduced Boyer’s costs by \$26/ac, and without any loss in corn yield, those reduced costs directly translated to \$26/ac in improved economic returns.

Boyer has been using a cereal rye cover crop in his corn-soybean rotation because he knows this practice is beneficial to the soil and reduces loss of nutrients, like N, to the environment. After six years of using a cereal rye cover crop and capturing those environmental benefits on his family’s farm, it now appears that he is also capturing economic benefits as well owing to reduced N fertilizer costs. Encouraged by these results, Boyer wonders if he can further reduce his N fertilizer rate and improve returns.

APPENDIX – TRIAL DESIGN AND WEATHER CONDITIONS

130 lb N/ac	180 lb N/ac	130 lb N/ac	180 lb N/ac	130 lb N/ac	180 lb N/ac	130 lb N/ac	180 lb N/ac
REP 1		REP 2		REP 3		REP 4	

FIGURE A1. Jack Boyer’s experimental design. He implemented four replications of the two N rate treatments (eight strips total). This design allows for statistical analysis of the results.

Grundy Center




FIGURE A2. Mean monthly temperature and rainfall for Apr. 1, 2019 through Oct. 31, 2019 and the long-term averages at Grundy Center, the nearest weather station to Boyer’s farm (about 9 miles away).^[4]

FUNDING ACKNOWLEDGEMENT

This material is based upon work supported by the Natural Resources Conservation Service, U.S. Department of Agriculture, under number 68-6114-17-010. Any opinions, findings, conclusions, or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the views of the U.S. Department of Agriculture.

REFERENCES

1. Sawyer, J.E. and A.P. Mallarino. 2017. Use of the Late-Spring Soil Nitrate Test in Iowa Corn Production. CROP 3140 Iowa State University Extension and Outreach. <https://store.extension.iastate.edu/Product/5259> (accessed October 2019).
2. Sawyer, J. and A.P. Mallarino. 2018. Cornstalk Testing to Evaluate Nitrogen Management. PM 1584. Iowa State University Extension and Outreach. <https://store.extension.iastate.edu/product/Use-of-the-End-of-Season-Corn-Stalk-Nitrate-Test-in-Iowa-Corn-Production> (accessed October 2019).
3. Plastina, A., A. Johanns and G. Wynne. 2019. 2019 Iowa Farm Custom Rate Survey. Ag Decision Maker. Iowa State University Extension and Outreach. <https://store.extension.iastate.edu/product/2019-Iowa-Farm-Custom-Rate-Survey> (accessed November 2019).
4. Iowa Environmental Mesonet. 2019. Climodat Reports. Iowa State University. <http://mesonet.agron.iastate.edu/climodat/> (accessed November 2019).



PFI COOPERATORS’ PROGRAM

PFI’s Cooperators’ Program helps farmers find practical answers and make informed decisions through on-farm research projects.

The Cooperators’ Program began in 1987 with farmers looking to save money through more judicious use of inputs.

If you are interested in conducting an on-farm trial contact Stefan Gailans @ 515-232-5661 or stefan@practicalfarmers.org.