



## Cauliflower Variety Trial

### In a Nutshell:

- Mark Quee and Shanti Sellz evaluated yield of fall-harvested cauliflower varieties.

### Key Findings:

- Mardi performed best among all varieties trialed in Quee's first succession. No differences among varieties were observed in the second succession, though.
- Snow Crown was the only variety harvested at Sellz's.

### Cooperators

Mark Quee, Scattergood Friends Farm  
 (West Branch)  
 Shanti Sellz, Muddy Miss Farm  
 (Iowa City)

### Funding

Ceres Trust

### METHODS

#### Design

Quee and Sellz seeded cauliflower varieties in late spring and transplanted in mid-summer for fall harvest. Quee included two successions; Sellz included only one. Management at both farms is shown in **Table 1**.

Both cooperators planted cauliflower varieties in randomized and replicated strips (**Figure A1**).

TABLE 1. Management practices at both farms in 2019.

	QUEE	SELLZ
Seeding date(s)	Succession 1 (S1): Apr. 30 Succession 2 (S2): June 11	June 21
Transplant date(s)	S1: June 10; S2: July 2	Aug. 7
Harvest date(s)	Sept. 24 - Oct. 27	Oct. 24 - Nov. 8
Varieties	Amazing; Denali; Mardi; Skywalker; Snow Crown (S2 only)	Amazing; Denali; Skywalker; Snow Crown
In-row spacing	24 in.	18 in.
Between-row spacing	30 in.	36 in.
Irrigation	Drip	Pressurized drip
Mulch	N	N
Banding	N	N
Plants per plot	S1: 9; S2: 15	20



Snow Crown cauliflower at Shanti Sellz's.

#### Measurements

Cooperators harvested cauliflower from each individual plot. Upon harvest, cooperators recorded number of cauliflower curds per plot, individual curd diameter, individual curd weight and the total weight of all curds harvested from the plot.

#### Data analysis

To rank the performance of cauliflower varieties, we calculated Tukey's least significant difference (LSD) for each measurement: harvested curds, average curd diameter, average curd weight and total plot weight. For each measurement, we assigned different rankings to the varieties if the difference observed between any two varieties was greater than or equal to the LSD – we also refer to this as a statistically significant effect. On the other hand, if the difference observed between any two varieties was less than the LSD, we consider the varieties to be statistically similar (those two varieties received the same ranking). We used a 90% confidence

level to calculate the LSDs, which means that we would expect our rankings to occur 9 times out of 10. We could make these statistical calculations because the farmers' experimental design involved replication of the cauliflower varieties (**Figure A1**).

## RESULTS AND DISCUSSION

### Quee

Succession 1 was started and transplanted about 1 month earlier than succession 2, but both groups were last harvested on October 27. Mardi performed better than all other varieties in succession 1, producing more curds per plot and larger curds than any other variety (**Table 2**). Denali and Amazing were middle of the pack, with Skywalker having the smallest curds. No statistical differences among varieties were observed during succession 2, which had fewer plots reach maturity due to heavy bird predation on the plants.

### Sellz

Sellz was only able to harvest Snow Crown curds; no other varieties produced curds for her (**Table 3**). She had similar results during a 2018 trial, and wondered if it was related to the weather that year, or the field location. Of the varieties chosen, Snow Crown has the fewest days to maturity – 50 days compared to 80 days for Skywalker (68 for Amazing, 73 for Denali). Planting dates were appropriate for fall planting; the plants had 93 days in the field after transplanting. Numerous factors impact curd development including nutrition (cauliflower are heavy feeders), but most commonly, temperature – cauliflower curd development is even more sensitive to temperature than broccoli head development. A condition called “blind bud” can occur, where the plant will not form a central bud and subsequently, not form a head.



Cauliflower varieties in the field at Mark Quee's in September 2019.

TABLE 2. Cauliflower yiled components for both successions at Quee's in 2019.  
*Top yield performer highlighted.*

SUCCESSION	VARIETY	HARVESTED CURDS (no./plot)	CURD DIAMETER (in.)	CURD WEIGHT (lb)	PLOT WEIGHT (lb)
S1	Amazing	4.1 ab	5.5 ab	0.91 ab	3.8 b
	Denali	3.6 ab	4.6 bc	0.82 b	2.9 b
	Mardi	5.6 a	6.4 a	1.48 a	8.4 a
	Skywalker	1.6 b	4.2 c	0.60 b	1.3 b
	LSD	2.8	1.11	0.58	4.2
S2	Amazing	1.8	5.0	1.01	1.5
	Denali	1.9	4.6	0.68	1.3
	Mardi	2.3	4.6	0.85	2.0
	Skywalker	0.3	5.0	0.88	0.2
	Snow Crown	2.8	5.7	1.34	3.1
LSD	3.7	1.6	1.17	3.2	

In succession 1 (S1), results that differ by more than the least significant difference (LSD) are followed by a different letter-ranking and are considered statistically different at the 90% confidence level. In succession 2 (S2), no letters appear because no results differed by more than the LSD; results are thus considered statistically similar among varieties.

Sellz noted that the transplants were in pretty rough shape when they went into the field due to being late with transplanting and irregular watering. Stressed transplants could have led to the later issues with curd development, even though the plants survived.

TABLE 3. Cauliflower yield components at Sellz's in 2019.

VARIETY	HARVESTED CURDS (no./plot)	CURD DIAMETER (in.)	CURD WEIGHT (lb)	PLOT WEIGHT (lb)
Snow Crown	3.25	3.0	0.24	0.77

Snow Crown was the only variety that produced harvestable curds.

## CONCLUSIONS AND NEXT STEPS

"I had never grown cauliflower in Iowa before conducting this trial," Sellz said. "From my experience and practices, Snow Crown will be my go to variety for future fall plantings."

"Amazing and Denali have been my go-to varieties for summer production," Quee said. "I'll have to look harder at Mardi. I didn't have any prior experience with Mardi and will definitely continue to see how it fits in my successions." Going forward, he plans on using all of these varieties except for Skywalker. "Amazing, Denali, Mardi and Snow Crown provide a nice harvest window from a single planting/transplanting and seem to account for a lot of variables."

## APPENDIX – TRIAL DESIGN AND WEATHER CONDITIONS

REP 1	REP 2	REP 3	REP 4
Mardi	Amazing	Denali	Mardi
Amazing	Skywalker	Mardi	Amazing
Skywalker	Denali	Amazing	Skywalker
Denali	Mardi	Skywalker	Denali

FIGURE A1. Example experimental design: four replications of four cauliflower varieties (16 plots total). This design allows for statistical analysis of the results.

TABLE A1. Climate data for 2019 and historical averages from the nearest weather station to both farms (Iowa City).

MONTH	GROWING DEGREE DAYS (base 50 °F)		RAINFALL (in.)	
	2019	Avg.	2019	Avg.
May	344	430	<b>9.6</b>	<b>4.4</b>
June	621	646	3.4	5.1
July	843	753	<b>1.0</b>	<b>4.0</b>
August	711	699	4.7	4.3
September	629	486	<b>7.5</b>	<b>3.1</b>
October	181	260	2.1	2.9
November	20	66	1.6	2.2

Climate data were accessed from the Iowa City weather station.<sup>[1]</sup> Historical data include years 1985-2018. Where rainfall in 2019 was more than two inches different than the average, values are displayed in **bold**.

## REFERENCES

1. Iowa Environmental Mesonet. 2020. Climodat Reports. Iowa State University. <http://mesonet.agron.iastate.edu/climodat/> (accessed October 2020).



### **PFI COOPERATORS' PROGRAM**

PFI's Cooperators' Program helps farmers find practical answers and make informed decisions through on-farm research projects.

The Cooperators' Program began in 1987 with farmers looking to save money through more judicious use of inputs.

If you are interested in conducting an on-farm trial contact Stefan Gailans @ 515-232-5661 or [stefan@practicalfarmers.org](mailto:stefan@practicalfarmers.org).